

CONSULTING

MURDOCH

Suite 27, Level 1
Murdoch Medical Clinic
St John of God Hospital
100 Murdoch Drive
Murdoch
WA 6150

WEST LEEDERVILLE

162 Cambridge Street
West Leederville
WA 6007

BOORAGOON

Garden City Specialist Centre
537 Marmion Street
Booragoon
WA 6154

MANDURAH

117 Anstruther Road
Mandurah
WA 6210

OPERATING

St John of God Hospital
Murdoch

St John of God Hospital
Subiaco

Mount Hospital
Perth

For all appointments

Phone: (08) 9332 0066

Fax: (08) 9312 1619 (Admin)
(08) 9463 6202 (Clinical)

**We Prefer Health Link
EDI - prthsurg**



REVIEW ARTICLE

OPEN ACCESS

Obesity and menopause

Santiago Palacios^a , Peter Chedraui^b , Rafael Sánchez-Borrego^c , Pluvio Coronado^d and Rossella E. Nappi^e

The absence of oestrogens, a general mechanism for fat gain operates by reducing the metabolic rate at rest, decreasing spontaneous physical activity, and increasing caloric intake – even with HRT – but HRT does seem to preserve muscle mass.

Weight may not change at menopause, but fat may increase and lean body mass decrease.

Resting energy expenditure is lower during the post menopause than in the pre-menopause

During female midlife, obesity is associated with menstrual cycle alterations (e.g. anovulatory cycles ending with abnormal uterine bleeding), menopausal symptoms including hot flashes, poor sleep, aches and joint pain, urinary symptoms, and reduced quality of life

While Bone Density is higher in obesity, it may not be sufficiently increased to compensate for the negative factors affecting bone health, including biomechanical factors such as bone geometry and quality. This may be related to an increase falls risk in obesity.

Women with overweight or obesity are more likely to experience urinary incontinence. A recent meta-analysis reveals a strong association between high BMI, abdominal pressure, and bladder pressure

Obesity also increases the severity of symptoms of impaired sexual functioning, unbalancing the levels of sex hormones, hence, resulting in a reduction of sexual desire, arousal, and orgasm


As BMI increases, the likelihood of urinary incontinence and shortness of breath also increases

Scientific societies agree on the fact of considering MHT as safe and a recommendable management option in symptomatic menopausal women under the age of 60 or less than 10 years after menopause. Although the benefits of MHT in these women have been demonstrated, by reducing cardiovascular risk, osteoporosis, and bone fractures, while also improving genitourinary syndrome of menopause, and quality of life, MHT should not be solely prescribed, as an isolated indication, for preventive purposes, except in case of premature ovarian insufficiency.

Women with increased BMI have a higher risk of thromboembolic disease (TED) when using MHT. But the actual absolute increase in risk is low. In women less than 60 and should not be a reason not to prescribe.

The use of oestrogen only or combined transdermal MHT does not increase the risk of thrombotic events in women with obesity

Summary by A/Prof HS Chandraratna



Obesity and menopause

Santiago Palacios, Peter Chedraui, Rafael Sánchez-Borrego, Pluvio Coronado & Rossella E. Nappi

To cite this article: Santiago Palacios, Peter Chedraui, Rafael Sánchez-Borrego, Pluvio Coronado & Rossella E. Nappi (2024) Obesity and menopause, Gynecological Endocrinology, 40:1, 2312885, DOI: [10.1080/09513590.2024.2312885](https://doi.org/10.1080/09513590.2024.2312885)

To link to this article: <https://doi.org/10.1080/09513590.2024.2312885>



© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 11 Feb 2024.



Submit your article to this journal [↗](#)



Article views: 2665








View related articles [↗](#)



View Crossmark data [↗](#)

Obesity and menopause

Santiago Palacios^a , Peter Chedraui^b , Rafael Sánchez-Borrego^c , Pluvio Coronado^d  and Rossella E. Nappi^e 

^aPalacios's Clinic for Women's Health, Madrid, Spain; ^bEscuela de Posgrado en Salud, Universidad Espíritu Santo, Samborondón, Ecuador; ^cDiatros, Woman's Clinic, Barcelona, Spain; ^dWomen's Health Institute, Hospital Clínico San Carlos, IdISSC, School of Medicine, Complutense University, Madrid, Spain; ^eDepartment of Clinical, Surgical, Diagnostic and Pediatric Sciences, University of Pavia, Research Center for Reproductive Medicine, Gynecological Endocrinology and Menopause, IRCCS San Matteo Foundation, Pavia, Italy

ABSTRACT

Obesity is not a choice or a result of lack of willpower, but a multifactorial, chronic, progressive, and relapsing disease. During menopause, hormonal and body composition changes lead to greater visceral adiposity, that aggravates women's health at a cardiometabolic, mechanic and mental level. Adiposity has been identified as an important modifier of reproductive hormones. During female midlife, obesity has been associated with menstrual cycle alterations (anovulatory cycles ending with abnormal bleedings), menopausal symptoms including hot flashes, poor quality of sleep, aches and joint pain, genitourinary symptoms, and reduced quality of life. However, the relationships between weight, the menopausal process, aging, and hormone levels remain poorly understood. Women with obesity have an increased risk of thromboembolic disease when using menopause hormone therapy (MHT), and it is probably the main medical condition to prescribe or not MHT. However, this risk depends on the route and type of MHT. The use of estrogen-only or combined transdermal MHT does not increase the risk of a thrombotic event in women with obesity.

ARTICLE HISTORY

Received 12 December 2023
Revised 3 January 2024
Accepted 25 January 2024
Published online 12 February 2024

KEYWORDS

Obesity; menopause; vasomotor symptoms; fracture risk; hormone therapy

Introduction

Obesity is not a 'choice', it arises from a complex interplay of biological, psychological, environmental and social factors. Obesity is not simply related to an individual's lack of willpower [1], rather, it is a chronic, progressive, and relapsing disease that affects women at higher rates than men [2]. According to the World Health Organization (WHO), overweight and obesity are defined as the abnormal or excessive accumulation of fat, that imposes risks to health, quality of life and life expectancy [3]. Excess of ectopic body fat can lead to the production of adipocytokines and inflammatory mediators, that disrupt glucose and fat metabolism, thus, increasing cardiometabolic and cancer risks [4].

In a gynecological setting, complications frequently encountered related to obesity include menopause problems/symptoms [5] and the increased risk of breast and gynecological cancers [6]. The endocrine tumult of the menopausal transition also exposes racial and socioeconomic disparities regarding the onset, severity, and frequency of symptoms [5,6]. There is ever-growing evidence of the impact that the excess of adiposity observed during the menopause and after, mainly due to biological changes resulting from decreased estrogen levels, has on various health indicators [7]. On the other hand, obesity can influence the decision to prescribe menopausal hormone therapy (MHT) [5,7].

Literature search

For this narrative review we performed a literature search using PubMed®, Embase®, and Scopus® to cover all publications up

to November 2023. The search terms included the following keywords in the title and/or the abstract: obesity, menopause, fat mass, hormone therapy, vasomotor symptoms, osteoporosis. The search included studies in English language. A manual check was performed for potential additional studies.

How does the menopause affect fat mass?

Studies suggest that both estradiol (E2) and follicle-stimulating hormone (FSH) play roles in regulating energy balance, and hence, their known variations during the menopausal transition may influence fat mass quantity and composition [8,9]. E2 affects numerous energy homeostasis pathways, such as central nervous system (CNS) control of food intake and energy expenditure, regulation of lipid storage and metabolism in the adipose tissue, and insulin sensitivity [8]. Experiments involving laboratory animals (e.g. estrogen receptor Knockout and Knock-in models and ovariectomy with and without hormone supplementation) have shown that, in the absence of estrogens, a general mechanism for fat gain operates by reducing the metabolic rate at rest, decreasing spontaneous physical activity, and increasing caloric intake [10].

Several cross-sectional and longitudinal observational studies have found that resting energy expenditure (REE) is lower during the postmenopause than in the premenopause [11,12]. In premenopausal women, pharmacologic suppression of sex hormones through sustained administration of a gonadotropin-releasing hormone agonist (GnRH-a) reduces REE, whereas adding back

transdermal E2 compensates for this GnRH-a induced decrease [13]. Notably, this paradigm of pharmacological hormone suppression, with and without the addition of transdermal E2, results in loss of lean mass, as assessed by Dual energy X-ray absorptiometry (DXA) only in women not receiving E2 treatment [14].

In a sample of 130 women of the Women's Health Across the Nation (SWAN) study showed no effect of the time since the last menstrual period over fat mass or lean mass (as determined by bioelectrical impedance). Indeed, the trend showed a linear increase in fat mass and a slight linear decrease in lean mass over time [15]. Similar results were corroborated in another study involving 191 healthy women (35 to 45 years) who were evaluated every 5 years and 75 women (46 or more years) who were still menstruating and evaluated every 6 months [16]. Most existing studies that examine the relationship between advancement of the menopausal stage based on menstrual pattern (i.e. pre-, peri-, or postmenopause) and changes in body composition or weight have concluded that the menopause had no effect on these parameters [17–24]. Although weight increased over time, in most studies [17–22], they often have important limitations such as the lack of details within such broad periods of pre-, peri- and postmenopause and in some cases had long periods between evaluations [16–21].

After menopause, there is a clear increase in fat mass, and a decrease in lean mass. This likely explains why there is no observable accelerated increase in weight or body mass index (BMI) across the menopausal transition. Data are consistent with the growing observation that, while BMI is a well-established indicator of cardiometabolic risk, it is not nearly as informative as the adiposity index. Specific aspects of adiposity, such as fat distribution, are not detected by BMI [25,26], making this index a less useful indicator of cardiometabolic risk in older women [27]. Despite this, recently more value, as a risk predictor, has been given to the abdominal circumference measurement [28].

Ultimately, the menopausal transition is associated with accelerated gain of fat mass and the simultaneous loss of lean mass; but their combined rates of change do not result in a detectable acceleration in weight or BMI gain at the onset of the menopausal transition [29].

How does obesity/overweight affect menopause?

Adiposity has been identified as an important modifier of reproductive hormones. During female midlife, obesity is associated with menstrual cycle alterations (e.g. anovulatory cycles ending with abnormal uterine bleeding), menopausal symptoms including hot flashes, poor sleep, aches and joint pain, urinary symptoms, and reduced quality of life [30]. However, the relationship between weight, the menopausal process, aging, and hormone levels remains poorly understood [31].

In a longitudinal study conducted in the population-based Penn Ovarian Aging Cohort ($n=436$), which evaluated during 12 years anthropometric measures, menopausal status, and reproductive hormone quantifications [30], the findings clearly indicate that obesity is an important factor influencing the hormonal changes observed during the menopausal transition independent of age, race, and smoking.

Effect of obesity on age at menopause

Obesity at different ages (e.g. early versus mid-adulthood) may have different effects on the timing of menopause [32], due to

changes in the reproductive function over time; however, few studies have directly and closely examined this. Furthermore, weight change and weight regain may be associated with menopause timing, especially in the context of overall adiposity.

Current evidence on the association between BMI and age at menopause remains unclear. The SWAN found no association between obesity and the age at natural menopause, although an association was found between surgical menopause and obesity [33]. However, an international pooled analysis of 11 prospective studies [34] supported a previously reported association between higher BMI and later menopause.

A prospective cohort study examined a variety of adiposity measures in 78,759 premenopausal women from the Nurses' Health Study who were followed from 1989 to 2011 in order to determine the incidence of early natural menopause [35]. The authors observed a non-linear, J-shaped association between BMI and the risk of early natural menopause. Compared to lean-normal weight women with a BMI of 18.5–22.4 kg/m², those with underweight significantly had 30% higher odds of early menopause. In contrast, women with overweight had a significant 21–30% lower odds. Intermediate abdominal adiposity was associated with a lower risk when compared to the lowest level, which persisted after BMI adjustment. The heterogeneity of study populations, assessment methods, and definitions of obesity may explain the observed discrepancies between findings.

Interestingly, according to a systematic review and meta-analysis [36], ovarian reserve markers, Anti-Müllerian hormone (AMH) and FSH, were significantly lower in women with obesity than in those without obesity, and BMI was negatively correlated with AMH levels in all study populations. Beyond its value as a marker of ovarian reserve and potential for ovulation, it has been suggested that lower AMH serum levels during the menopausal transition could be associated with higher markers of obesity and may be predictive of future obesity-related cardiometabolic complications [37].

Effect of obesity on vasomotor symptoms

The association between BMI and vasomotor symptoms (VMS; hot flashes and night sweats) remains highly controversial. Menopause-induced changes in thermoregulatory neurons result in a narrowing of the thermoneutral zone which is coupled with a reduction of the temperature threshold for a compensatory vascular response [38]. To date, two contradictory hypotheses have been proposed to explain the mechanisms that determine the relation between BMI and VMS. The 'thermoregulatory model' [32,33] that suggests that higher BMI and adiposity are linked to a greater prevalence of hot flashes due to body adipose tissue acting as a thermal insulator, hindering heat dissipation, thus women with obesity would suffer more VMS [39,40]; on the other hand there is the so-called 'thin hypothesis' that proposes that women with excess body weight experience fewer VMS because the cytochrome P450 aromatase enzyme in fat tissue converts androgens to estrogens [41,42]. In this second model, body fat would serve as a supplementary source of estrogen by aromatizing androstenedione, converting it into estrone [43].

Similarly, the effect of obesity on menopause-related symptoms may differ in relation to age (early versus mid-adulthood) [33]; higher BMI is significantly associated with more hot flashes during the menopausal transition, whereas it would be linked to a lower risk of hot flashes in the postmenopause [44], where a higher level of estrone might reduce hot flash frequency [42].

Recent studies have explored the role of adipose tissue-derived cytokine-like substances, collectively known as adipokines, as an additional mechanism that might potentially explain the link between adiposity and flushing. Adiponectin, the most abundant adipokine in the body, which is anti-inflammatory, has been associated with improved cardiovascular health, including a lower risk of cardiovascular disease (CVD) events [45]. In women with obesity, low circulating adiponectin levels, as well as central resistance to leptin, may contribute to a greater severity of menopause-related symptoms [44]. High leptin levels and low adiponectin levels have been observed in premenopausal and perimenopausal women [38], as well as in postmenopausal women with hot flashes compared to those without hot flashes [46]. However, a cross-sectional analysis of 898 postmenopausal women found an association between self-reported VMS and higher levels of ghrelin, the orexigenic peptide hormone released by endocrine cells in the stomach [47], but not adiponectin [44]. Therefore, although there appears to be links between VMS and adiponectin [48], this association remains underexplored.

Some research seems to point out at the link between VMS and adverse adipokine profiles or cytokines produced by the adipose tissue. Physiologic VMS have been associated with lower adiponectin levels, after considering potential confounders, indicating that the potential role of adipokines in producing VMS and their links to health warrant further attention [46]. Understanding the relationship between menopausal symptoms and obesity may help identify new strategies to prevent weight gain during female midlife, and thus, reduce adverse outcomes related to obesity and the severity of VMS [49].

Obesity, osteoporosis, and fracture risk: the 'obesity paradigm'

The belief that obesity is protective against fractures has influenced clinical practice, with BMI being part of the fracture risk assessment tool (FRAX). Studies have shown a positive correlation between bone mineral density (BMD) and BMI [50] and a lower incidence of hip fractures in adults with obesity [51]. However, accumulating evidence challenges this perception, which is known as the 'obesity paradigm'. While BMD is higher in obesity [52], it may not be sufficiently increased to compensate for the negative factors affecting bone health, including biomechanical factors such as bone geometry and quality.

A systematic review and meta-analysis [53] of 121 studies reported higher lumbar spine, total hip, femoral neck, and radius BMD among men and women (pre- and postmenopausal) with obesity compared to their counterparts with normal weight. Contrary to this, a recent retrospective study [54] identified abdominal obesity as a negative predictor of bone health among older adults, independent of BMI, with an inverted U-shaped association, and a point of inflection at a waist circumference of 95 cm, considered as a clinical parameter of abdominal obesity.

Although the underlying mechanisms linking obesity and bone health remain unknown, the common origin of osteoblasts and adipocytes from stromal cells may be the explanation of the link between adiposity and bone health [55]. The coexistence of a higher BMD and a higher prevalence of fractures suggests that in these patients, bone density is either more fragile or they are more prone to falls. Both could be partly explained by the consequences of impaired glucose metabolism. Changes in the properties of bone collagen material, due to the accumulation of advanced glycation end-products have been described [56]. Chances of falling are increased in patients with diabetes due to

polyneuritis or episodic-hypoglycemia. Over time, it can be assumed that the accumulation of advanced glycation end products, as well as vascular and neural complications, may contribute to a greater tendency to fracture instead of offering protection against fractures due to greater bone mass [57].

Effect of obesity on other specific parameters of the menopause

Obesity has been associated with several other specific menopausal parameters or aspects including urinary symptoms, sexual functioning and quality of life. Women with overweight or obesity are more likely to experience urinary incontinence (UI) substantially affecting their quality of life [58,59]. Overweight and obesity have been identified as an important predictor of UI, with the risk being significantly elevated in cases of obesity. A recent meta-analysis reveals a strong association between high BMI, abdominal pressure, and bladder pressure [60]. Therefore, currently, obesity and overweight are considered independent risk factors for UI [59,60].

An important clinical condition to consider in postmenopausal women is the presence of sarcopenic obesity defined by excess adiposity and low skeletal muscle mass and/or function [61]. Sarcopenic obesity is a predictor of disability and survival, dramatically increasing the risk of obesity and age-related diseases [61].

Consistent evidence shows a bidirectional link between overweight/obesity and emotional distress in the general population [62]. In fact, adipokines produced by the adipose tissue activate systemic inflammation, which eventually affects the brain causing mood dysregulation [63]. It has also been hypothesized that menopause-related mood disorders may lead to increased cytokine and free radical generation, contributing to increased fat accumulation [64].

Furthermore, obesity also increases the severity of symptoms of impaired sexual functioning, unbalancing the levels of sex hormones, hence, resulting in a reduction of sexual desire, arousal, and orgasm [65]. Additionally, as obesity favors the likelihood of developing comorbidities such as type 2 diabetes, dyslipidemia and hypertension, these conditions in turn increase the risk of reduced sexual activity [66]. Moreover, people with obesity are more likely to experience anxiety and/or depression, which are thought to directly or indirectly impact sexual function [65,67]. As BMI increases, the likelihood of urinary incontinence and shortness of breath also increases [68]; however, in contrast, the relationship between BMI and the likelihood of aches/stiffness is nonlinear, increasing to a threshold level [68].

Risks and benefits of menopause hormone therapy in women living with obesity

MHT has been indicated in women suffering from menopausal symptoms; however, a high proportion of eligible women (54%–79%) are unwilling to receive this option [69]. According to recent guidelines [70–72] MHT remains the primary treatment for menopausal symptoms, particularly VMS and this statement has not changed in the last few years [73]. As it is recommended for medical interventions, MHT should be individualized according to women's needs, symptoms, and clinical conditions. In addition, it is relevant to provide the best MHT in terms of type, route, dose, and duration considering efficacy, tolerability, adherence, and any other relevant aspects [72]. In this sense, obesity

should be kept in mind when prescribing MHT, because adipose tissue is an endocrine tissue that produces estrogens [74] and it is associated with other disorders, such as the metabolic syndrome [75].

Scientific societies agree on the fact of considering MHT as safe and a recommendable management option in symptomatic menopausal women under the age of 60 or less than 10 years after menopause [71,72]. Although the benefits of MHT in these women have been demonstrated, by reducing cardiovascular risk, osteoporosis, and bone fractures, while also improving genitourinary syndrome of menopause, and quality of life [76,77], MHT should not be solely prescribed, as an isolated indication, for preventive purposes [77], except in case of premature ovarian insufficiency [78].

Women with increased BMI have a higher risk of thromboembolic disease (TED) when using MHT (OR 2.5, 95% CI, 1.7-3.7 [overweight]; OR 3.9, 95% CI, 2.2-6.9 [obesity]) as compared to women with normal weight [79]. TED is probably the main medical condition to prescribe or not MHT. However, this risk depends on the route and type of MHT and the severity of obesity. Women with overweight face an increased risk of TED when using oral MHT (combined or estrogen-only), although the absolute risk, especially in women under the age of 60 is low [80]. Transdermal MHT (with or without progesterone), on the other hand, is not associated with an increased risk; hence, the transdermal route should be preferred in patients with a risk or history of TED [80].

There are few studies that analyze the risk of using any type of MHT in obese women (BMI \geq 30 kg/m²). This is the reason why currently the prescription of MHT must be individualized, taking into consideration risks and benefits. Despite this, data regarding transdermal MHT suggest that the use of estrogen-only or combined transdermal MHT does not increase the risk of a thrombotic event in women with obesity (BMI \geq 30 kg/m² but <35 kg/m²) [81,82].

Conclusions

During the menopausal transition there are unfavorable changes in body composition that include an increase in fat mass, particularly in the abdominal region, and a decrease in fat-free mass [31,83]. Whether these modifications are primarily due to age or hormonal changes is still a matter of intensive research and debate [31,83,84].

We currently consider that BMI is an independent factor for the presence of hot flashes, along with urge incontinence and vaginal dryness, supporting the thermoregulatory theory [85]. While BMD is higher in obesity [52], it may not be sufficiently increased to compensate for negative factors that affect bone health, including biomedical and biomechanical factors such as bone geometry and quality. For this reason, our belief is that all these factors predispose a greater tendency for fractures instead of offering protection against them due to greater bone mass.

In women with obesity, combined oral MHT should not be the first choice because the evidence shows an increased risk of TED; although the absolute risk, especially in those under 60 years of age, is low. The use of estrogen-only or combined transdermal MHT does not increase the risk of thrombotic events in women with obesity [81].

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

The author(s) reported there is no funding associated with the work featured in this article.

ORCID

Santiago Palacios  <http://orcid.org/0000-0003-2229-1200>

Peter Chedraui  <http://orcid.org/0000-0002-1556-3979>

Rafael Sánchez-Borrego  <http://orcid.org/0000-0001-6583-7440>

Pluvio Coronado  <http://orcid.org/0000-0003-0357-2015>

Rossella E. Nappi  <http://orcid.org/0000-0002-5216-9882>

Data availability statement

Data sharing not applicable – no new data generated.

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

References

- [1] Chater A. Obesity. Psychological perspectives on obesity: policy, practice and research priorities. The British Psychological Society. 2020 May 7. Available from: <https://www.bps.org.uk/blog/psychological-perspectives-obesity-policy-practice-and-research-priorities>
- [2] Leeners B, Geary N, Tobler PN, et al. Ovarian hormones and obesity. Hum Reprod Update. 2017;23(3):1–7. doi:10.1093/humupd/dmw045.
- [3] World Health Organization. Obesity and overweight. Available from: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
- [4] Wharton S, Lau DCW, Vallis M, et al. Obesity in adults: a clinical practice guideline. CMAJ. 2020;192(31):E875–E891. doi:10.1503/cmaj.191707.
- [5] Palacios S. Obesity in women's life: role of GLP-1 agonists. Gynecol Endocrinol. 2022;38(11):889–890. doi:10.1080/09513590.2022.2163235.
- [6] Wichmann IA, Cuello MA. Obesity and gynecological cancers: a toxic relationship. Int J Gynaecol Obstet. 2021;155(Suppl 1):123–134. doi:10.1002/ijgo.13870.
- [7] Nappi RE, Chedraui P, Lambrinoudaki I, et al. Menopause: a cardiometabolic transition. Lancet Diabetes Endocrinol. 2022;10(6):442–456. doi:10.1016/S2213-8587(22)00076-6.
- [8] Mauvais-Jarvis F, Clegg DJ, Hevener AL. The role of estrogens in control of energy balance and glucose homeostasis. Endocr Rev. 2013;34(3):309–338. doi:10.1210/er.2012-1055.
- [9] Kohrt WM, Wierman ME. Preventing fat gain by blocking Follicle-Stimulating hormone. N Engl J Med. 2017;377(3):293–295. doi:10.1056/NEJMcibr1704542.
- [10] Van Pelt RE, Gavin KM, Kohrt WM. Regulation of body composition and bioenergetics by estrogens. Endocrinol Metab Clin North Am. 2015;44(3):663–676. doi:10.1016/j.ecl.2015.05.011.
- [11] Hodson L, Harnden K, Banerjee R, et al. Lower resting and total energy expenditure in postmenopausal compared with premenopausal women matched for abdominal obesity. J Nutr Sci. 2014;3:e3. doi:10.1017/jns.2013.38.
- [12] Lovejoy JC, Champagne CM, de Jonge L, et al. Increased visceral fat and decreased energy expenditure during the menopausal transition. Int J Obes (Lond). 2008;32(6):949–958. doi:10.1038/ijo.2008.25.
- [13] Melanson EL, Gavin KM, Shea KL, et al. Regulation of energy expenditure by estradiol in premenopausal women. J Appl Physiol (1985). 2015;119(9):975–981. doi:10.1152/jappphysiol.00473.2015.
- [14] Shea KL, Gavin KM, Melanson EL, et al. Body composition and bone mineral density after ovarian hormone suppression with or without estradiol treatment. Menopause. 2015;22(10):1045–1052. doi:10.1097/GME.0000000000000430.
- [15] Sowers M, Zheng H, Tomey K, et al. Changes in body composition in women over six years at midlife: ovarian and chronological aging. J Clin Endocrinol Metab. 2007;92(3):895–901. doi:10.1210/jc.2006-1393.
- [16] Davies KM, Heaney RP, Recker RR, et al. Hormones, weight change and menopause. Int J Obes Relat Metab Disord. 2001;25(6):874–879. doi:10.1038/sj.ijo.0801593.

- [17] Guthrie JR, Dennerstein L, Dudley EC. Weight gain and the menopause: a 5-year prospective study. *Climacteric*. 1999;2(3):205–211. doi:10.3109/13697139909038063.
- [18] Wing RR, Matthews KA, Kuller LH, et al. Weight gain at the time of menopause. *Arch Intern Med*. 1991;151(1):97–102. doi:10.1001/archinte.1991.00400010111016.
- [19] Crawford SL, Casey VA, Avis NE, et al. A longitudinal study of weight and the menopause transition: results from the Massachusetts women's health study. *Menopause*. 2000;7(2):96–104. doi:10.1097/00042192-200007020-00005.
- [20] Blümel JE, Castelo-Branco C, Rocangliolo ME, et al. Changes in body mass index around menopause: a population study of Chilean woman. *Menopause*. 2001;8(4):239–244. doi:10.1097/00042192-200107000-00004.
- [21] Sternfeld B, Wang H, Quesenberry CP, Jr, et al. Physical activity and changes in weight and waist circumference in midlife women: findings from the study of women's health across the nation. *Am J Epidemiol*. 2004;160(9):912–922. doi:10.1093/aje/kwh299.
- [22] Ho SC, Wu S, Chan SG, et al. Menopausal transition and changes of body composition: a prospective study in chinese perimenopausal women. *Int J Obes (Lond)*. 2010;34(8):1265–1274. doi:10.1038/ijo.2010.33.
- [23] Sornay-Rendu E, Karras-Guillibert C, Munoz F, et al. Age determines longitudinal changes in body composition better than menopausal and bone status: the OFELY study. *J Bone Miner Res*. 2012;27(3):628–636. doi:10.1002/jbmr.1469.
- [24] Abdunour J, Doucet E, Brochu M, et al. The effect of the menopausal transition on body composition and cardiometabolic risk factors: a Montreal-Ottawa new emerging team group study. *Menopause*. 2012;19(7):760–767. doi:10.1097/gme.0b013e318240f6f3.
- [25] Wells JC. Commentary: the paradox of body mass index in obesity assessment: not a good index of adiposity, but not a bad index of cardio-metabolic risk. *Int J Epidemiol*. 2014;43(3):672–674. doi:10.1093/ije/dyu060.
- [26] Müller MJ, Lagerpusch M, Enderle J, et al. Beyond the body mass index: tracking body composition in the pathogenesis of obesity and the metabolic syndrome. *Obes Rev*. 2012;13(Suppl 2):6–13. doi:10.1111/j.1467-789X.2012.01033.x.
- [27] Rubin R. Postmenopausal women with a "normal" BMI might be overweight or even obese. *JAMA*. 2018;319(12):1185–1187. doi:10.1001/jama.2018.0423.
- [28] Christiansen MR, Kilpeläinen TO, McCaffery JM. Abdominal obesity genetic variants predict waist circumference regain after weight loss. *Diabetes*. 2023;72(10):1424–1432. doi:10.2337/db23-0131.
- [29] Cienfuegos S, Gabel K, Kalam F, et al. Changes in body weight and metabolic risk during time restricted feeding in premenopausal versus postmenopausal women. *Exp Gerontol*. 2021;154:111545. doi:10.1016/j.exger.2021.111545.
- [30] Freeman EW, Sammel MD, Lin H, et al. Obesity and reproductive hormone levels in the transition to menopause. *Menopause*. 2010;17(4):718–726. doi:10.1097/gme.0b013e3181ccc85d.
- [31] Davis SR, Castelo-Branco C, Chedraui P, et al. Writing group of the international menopause society for world menopause day 2012. Understanding weight gain at menopause. *Climacteric*. 2012;15(5):419–429. doi:10.3109/13697137.2012.707385.
- [32] Thurston RC, Santoro N, Matthews KA. Adiposity and hot flashes in midlife women: a modifying role of age. *J Clin Endocrinol Metab*. 2011;96(10):E1588–1595. doi:10.1210/jc.2011-1082.
- [33] Thurston RC, Sowers MR, Chang Y, et al. Adiposity and reporting of vasomotor symptoms among midlife women: the study of women's health across the nation. *Am J Epidemiol*. 2008;167(1):78–85. doi:10.1093/aje/kwm244.
- [34] Zhu D, Chung HF, Pandeya N, et al. Body mass index and age at natural menopause: an international pooled analysis of 11 prospective studies. *Eur J Epidemiol*. 2018;33(8):699–710. doi:10.1007/s10654-018-0367-y.
- [35] Szegda KL, Whitcomb BW, Purdue-Smithe AC, et al. Adult adiposity and risk of early menopause. *Hum Reprod*. 2017;32(12):2522–2531. doi:10.1093/humrep/dex304.
- [36] Moslehi N, Shab-Bidar S, Ramezani Tehrani F, et al. Is ovarian reserve associated with body mass index and obesity in reproductive aged women? A meta-analysis. *Menopause*. 2018;25(9):1046–1055. doi:10.1097/GME.0000000000001116.
- [37] Francis EC, Oken E, Hivert MF, et al. Anti-Müllerian hormone and adiposity across mid-life among women in project viva. *Menopause*. 2023;30(3):247–253. doi:10.1097/GME.0000000000002143.
- [38] Navarro VM. Interactions between kisspeptins and neurokinin B. *Adv Exp Med Biol*. 2013;784:325–347. doi:10.1007/978-1-4614-6199-9_15.
- [39] Da Fonseca AM, Bagnoli VR, Souza MA, et al. Impact of age and body mass on the intensity of menopausal symptoms in 5968 Brazilian women. *Gynecol Endocrinol*. 2013;29(2):116–118. doi:10.3109/0951359.0.2012.730570.
- [40] Yokota M, Makita K, Hirasawa A, et al. Symptoms and effects of physical factors in Japanese middle-aged women. *Menopause*. 2016;23(9):974–983. doi:10.1097/GME.0000000000000660.
- [41] Thurston RC, Chang Y, Mancuso P, et al. Adipokines, adiposity, and vasomotor symptoms during the menopause transition: findings from the study of women's health across the nation. *Fertil Steril*. 2013;100(3):793–800. doi:10.1016/j.fertnstert.2013.05.005.
- [42] Gold EB, Crawford SL, Shelton JE, et al. Longitudinal analysis of changes in weight and waist circumference in relation to incident vasomotor symptoms: the study of women's health across the nation (SWAN). *Menopause*. 2017;24(1):9–26. doi:10.1097/GME.0000000000000723.
- [43] Anderson GS. Human morphology and temperature regulation. *Int J Biometeorol*. 1999;43(3):99–109. doi:10.1007/s004840050123.
- [44] Karim R, Dang HM, Hodis HN, et al. Association of hot flashes with ghrelin and adipokines in early versus late postmenopausal women. *Menopause*. 2020;27(5):512–518. doi:10.1097/GME.0000000000001508.
- [45] Hao G, Li W, Guo R, et al. Serum total adiponectin level and the risk of cardiovascular disease in general population: a meta-analysis of 17 prospective studies. *Atherosclerosis*. 2013;228(1):29–35. doi:10.1016/j.atherosclerosis.2013.02.018.
- [46] Huang WY, Chang CC, Chen DR, et al. Circulating leptin and adiponectin are associated with insulin resistance in healthy postmenopausal women with hot flashes. *PLoS One*. 2017;12(4):e0176430. doi:10.1371/journal.pone.0176430.
- [47] Kojima M, Hosoda H, Date Y, et al. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature*. 1999;402(6762):656–660. doi:10.1038/45230.
- [48] Thurston RC, Chang Y. Menopausal vasomotor symptoms and adiponectin among midlife women. *Menopause*. 2022;29(10):1145–1149. doi:10.1097/GME.0000000000002039.
- [49] Tang R, Fan Y, Luo M, et al. General and Central obesity are associated with increased severity of the VMS and sexual symptoms of menopause among chinese women: a longitudinal study. *Front Endocrinol (Lausanne)*. 2022;13:814872. doi:10.3389/fendo.2022.814872.
- [50] Gao B, Huang Q, Lin YS, et al. Dose-dependent effect of estrogen suppresses the osteo-adipogenic transdifferentiation of osteoblasts via canonical wnt signaling pathway. *PLoS One*. 2014;9(6):e99137. doi:10.1371/journal.pone.0099137.
- [51] Zhao LJ, Jiang H, Papiasian CJ, et al. Correlation of obesity and osteoporosis: effect of fat mass on the determination of osteoporosis. *J Bone Miner Res*. 2008;23(1):17–29. doi:10.1359/jbmr.070813.
- [52] Zhang Y, Pu J. The saturation effect of obesity on bone mineral density for older people: the NHANES 2017-2020. *Front Endocrinol (Lausanne)*. 2022;13:883862. doi:10.3389/fendo.2022.883862.
- [53] Turcotte AF, O'Connor S, Morin SN, et al. Association between obesity and risk of fracture, bone mineral density and bone quality in adults: a systematic review and meta-analysis. *PLoS One*. 2021;16(6):e0252487. doi:10.1371/journal.pone.0252487.
- [54] Chen J, Zhu L, Yao X, et al. The association between abdominal obesity and femoral neck bone mineral density in older adults. *J Orthop Surg Res*. 2023;18(1):171. doi:10.1186/s13018-023-03654-3.
- [55] Kume S, Kato S, Yamagishi S, et al. Advanced glycation end-products attenuate human mesenchymal stem cells and prevent cognate differentiation into adipose tissue, cartilage, and bone. *J Bone Miner Res*. 2005;20(9):1647–1658. doi:10.1359/JBMR.050514.
- [56] Gkataris K, Goulis DG, Potoupnis M, et al. Obesity, osteoporosis and bone metabolism. *J Musculoskelet Neuronal Interact*. 2020;20(3):372–381.

- [57] Yang Y, Hu X, Zhang Q, et al. Diabetes mellitus and risk of falls in older adults: a systematic review and meta-analysis. *Age Ageing*. 2016;45(6):761–767. doi:10.1093/ageing/afw140.
- [58] Yazdany T, Jakus-Waldman S, Jeppson PC, et al. American urogynecologic society. American urogynecologic society systematic review: the impact of weight loss intervention on lower urinary tract symptoms and urinary incontinence in overweight and obese women. *Female Pelvic Med Reconstr Surg*. 2020;26(1):16–29. doi:10.1097/SPV.0000000000000802.
- [59] Fuselier A, Hanberry J, Margaret Lovin J, et al. Obesity and stress urinary incontinence: impact on pathophysiology and treatment. *Curr Urol Rep*. 2018;19(1):10. doi:10.1007/s11934-018-0762-7.
- [60] Shang X, Fu Y, Jin X, et al. Association of overweight, obesity and risk of urinary incontinence in Middle-aged and older women: a meta epidemiology study. *Front Endocrinol (Lausanne)*. 2023;14:1220551. doi:10.3389/fendo.2023.1220551.
- [61] Gortan Cappellari G, Guillet C, Poggiogalle E, et al. Sarcopenic obesity research perspectives outlined by the sarcopenic obesity global leadership initiative (SOGLI) - Proceedings from the SOGLI consortium meeting in rome november 2022. *Clin Nutr*. 2023;42(5):687–699. doi:10.1016/j.clnu.2023.02.018.
- [62] Glaus J, Cui L, Hommer R, et al. Association between mood disorders and BMI/overweight using a family study approach. *J Affect Disord*. 2019;248:131–138. doi:10.1016/j.jad.2019.01.011.
- [63] Ali AM, Kunugi H. Royal jelly as an intelligent anti-Aging Agent-A focus on cognitive aging and Alzheimer's disease: a review. *Antioxidants (Basel)*. 2020;9(10):937. doi:10.3390/antiox9100937.
- [64] Elavsky S, Molenaar PCM, Gold CH, et al. Depressed mood but not fatigue mediate the relationship between physical activity and perceived stress in middle-aged women. *Maturitas*. 2009;71(3):287–293. doi:10.1016/j.maturitas.2011.12.011.
- [65] Abdelsamea GA, Amr M, Tolba AMN, et al. Impact of weight loss on sexual and psychological functions and quality of life in females with sexual dysfunction: a forgotten avenue. *Front Psychol*. 2023;14:1090256. doi:10.3389/fpsyg.2023.1090256.
- [66] Lyall DM, Celis-Morales C, Ward J, et al. Association of body mass index with cardiometabolic disease in the UK biobank: a Mendelian randomization study. *JAMA Cardiol*. 2017;2(8):882–889. doi:10.1001/jamacardio.2016.5804.
- [67] Rowland DL, McNabney SM, Mann AR. Sexual function, obesity, and weight loss in men and women. *Sex Med Rev*. 2017;5(3):323–338. doi:10.1016/j.sxmr.2017.03.006.
- [68] Sievert LL, Whitcomb BW, Verjee MA, et al. Limited evidence of a threshold effect for increasing adiposity on risk of symptoms at mid-life. *Menopause*. 2022;29(12):1381–1387. doi:10.1097/GME.0000000000002074.
- [69] Nappi RE, Kroll R, Siddiqui E, et al. Global cross-sectional survey of women with vasomotor symptoms associated with menopause: prevalence and quality of life burden. *Menopause*. 2021;28(8):875–882. doi:10.1097/GME.0000000000001793.
- [70] Baber RJ, Panay N, Fenton A. 2016 IMS recommendations on women's midlife health and menopause hormone therapy. *Climacteric*. 2016;19(2):109–150. doi:10.3109/13697137.2015.1129166.
- [71] Lambrinoudaki I, Armeni E, Goulis D, et al. Menopause, wellbeing and health: a care pathway from the european menopause and andropause society. *Maturitas*. 2022;163:1–14. doi:10.1016/j.maturitas.2022.04.008.
- [72] “The 2022 hormone therapy position statement of the North American menopause society” advisory panel. The 2022 hormone therapy position statement of the North American menopause society. *Menopause*. 2022;29(7):767–794.
- [73] Palacios S, Stevenson JC, Schaudig K, et al. Hormone therapy for first-line management of menopausal symptoms: practical recommendations. *Womens Health (Lond)*. 2019;15:1745506519864009. doi:10.1177/1745506519864009.
- [74] Bernasochi GB, Bell JR, Simpson ER, et al. Impact of estrogens on the regulation of white, beige, and brown adipose tissue depots. *Compr Physiol*. 2019;9(2):457–475.
- [75] Torres S, Fabersani E, Marquez A, et al. Adipose tissue inflammation and metabolic syndrome. The proactive role of probiotics. *Eur J Nutr*. 2019;58(1):27–43. doi:10.1007/s00394-018-1790-2.
- [76] Nappi RE. The 2022 hormone therapy position statement of the North American menopause society: no news is good news. *Lancet Diabetes Endocrinol*. 2022;10(12):832–834. doi:10.1016/S2213-8587(22)00285-6.
- [77] Trémollières FA, Chabbert-Buffet N, Plu-Bureau G, et al. Management of postmenopausal women: collège national des gynécologues et obstétriciens français (CNGOF) and groupe d'Etude sur la ménopause et le vieillissement (GEMVi) clinical practice guidelines. *Maturitas*. 2022;163:62–81. doi:10.1016/j.maturitas.2022.05.008.
- [78] Gartlehner G, Patel SV, Reddy S, et al. Hormone therapy for the primary prevention of chronic conditions in postmenopausal persons: updated evidence report and systematic review for the US preventive services task force. *JAMA*. 2022;328(17):1747–1765. doi:10.1001/jama.2022.18324.
- [79] Canonico M, Oger E, Conard J, et al. ESTrogen and THromboEmbolic risk (ESTHER) study group. Obesity and risk of venous thromboembolism among postmenopausal women: differential impact of hormone therapy by route of estrogen administration. The ESTHER study. *J Thromb Haemost*. 2006;4(6):1259–1265. doi:10.1111/j.1538-7836.2006.01933.x.
- [80] Mendoza N, Ramirez I, de la Viuda E, et al. Eligibility criteria for menopausal hormone therapy (MHT): a position statement from a consortium of scientific societies for the use of MHT in women with medical conditions. MHT eligibility criteria group. *Maturitas*. 2022;166:65–85. doi:10.1016/j.maturitas.2022.08.008.
- [81] Cushman M, Kuller LH, Prentice R, et al. Women's health initiative investigators. Estrogen plus progestin and risk of venous thrombosis. *JAMA*. 2004;292(13):1573–1580. doi:10.1001/jama.292.13.1573.
- [82] Vinogradova Y, Coupland C, Hippisley-Cox J. Use of hormone replacement therapy and risk of venous thromboembolism: nested case-control studies using the QResearch and CPRD databases. *BMJ*. 2019;364:k4810. doi:10.1136/bmj.k4810.
- [83] Wildman RP, Sowers MR. Adiposity and the menopausal transition. *Obstet Gynecol Clin North Am*. 2011;38(3):441–454. doi:10.1016/j.ogc.2011.05.003.
- [84] Al-Safi ZA, Polotsky AJ. Obesity and menopause. *Best Pract Res Clin Obstet Gynaecol*. 2015;29(4):548–553. doi:10.1016/j.bpobgyn.2014.12.002.
- [85] Saccomani S, Lui-Filho JF, Juliato CR, et al. Does obesity increase the risk of hot flashes among midlife women? A population-based study. *Menopause*. 2017;24(9):1065–1070. doi:10.1097/GME.0000000000000884.